



E-ISSN: 2789-1623
P-ISSN: 2789-1631
IJRP 2023; 3(1): 12-15
Received: 22-10-2022
Accepted: 26-11-2022

Tanul Jain
PG Resident, Department of
Psychiatry, Index Medical
College, Indore,
Madhya Pradesh, India

Dr. Ram Ghulam Razdan
Professor & Head, Department
of Psychiatry, Index Medical
College, Indore,
Madhya Pradesh, India

Study of prevalence of sexual dysfunction in males with psychiatric illness compared to other medical patients

Dr. Tanul Jain and Dr. Ram Ghulam Razdan

DOI: <https://doi.org/10.22271/27891623.2023.v3.i1a.36>

Abstract

Introduction: According to DSM-5 Sexual dysfunction is defined as “a clinically significant disturbance in a person's ability to respond sexually or to experience sexual pleasure”.

Aim & Objectives: To assess prevalence of sexual dysfunction in males with psychiatric illness compared to other medical patients in Psychiatry Department, Medicine Department, Respiratory Department of Index Medical College Hospital and Research Centre, and Indore.

Methodology: This study is a Comparative cross-sectional study conducted at Index Medical College Hospital and Research Centre, Indore for duration of 6 Months. Subjects was taken from department of Psychiatry, Medicine Department, Respiratory Department Index Medical College, Hospital and Research Centre, Indore fulfilling the inclusion criteria. Study sample will consist of 100 cases and 100 control.

Result: There was a positive relationship established between Psychiatric ill male patients and sexual dysfunction. Sexual dysfunction was significantly high in mentally ill patients i.e., 59% compared to our control group i.e. other medical patients (23%). Among the Psychiatric disorders highest frequency was observed in Schizophrenia which was 73%. Other prevalence was: bipolar disorders-49%, depressive disorders-51%, anxiety disorders-36% and Alcohol Use Disorder-58%. Among our control group the maximum frequency was observed in patients with heart diseases which was 41%. Other prevalence were liver disorders-25%, Diabetic-20%, genitourinary tract disorders-15% and resp. disorders-10%. It was also found that there was no significant difference in total testosterone level of both groups but prolactin level was significantly higher among psychiatric patients.

Conclusion: Prevalence of sexual dysfunctions are higher among psychiatric ill male patients than medical patients therefore more attention should be given to the treatment of these disorders and counselling should also be done.

Keywords: Sexual dysfunction, depression, anxiety

Introduction

Sexual dysfunction is defined as “a clinically significant disturbance in a person's ability to respond sexually or to experience sexual pleasure” (DSM-5). Sexual dysfunction is highly prevalent in Psychiatric patients which may be caused by the psychopathology of the illness or by the drugs given during their illness. Psychiatric disorders like Depression, Schizophrenia, Mood disorders, etc. may affect sexual relationship, sexual difficulties in patients. Find out the prevalence of sexual dysfunction among male psychiatric ill patients visiting Psychiatry OPD compared to other medical patients in Medicine Department & Respiratory Department of Index Medical College Hospital and Research Centre, and Indore.

Materials and Methods

This study is a comparative cross-sectional study conducted at Index Medical College Hospital and Research Centre, Indore for 6 Months (July to December 2022). Subjects was taken from department of Psychiatry, Medicine Department, Respiratory Medicine Department Index Medical College, Hospital and Research Centre, Indore fulfilling the inclusion criteria. Study sample will consist of 100 cases and 100 control. Study was done in two groups of male patients—group 1 consisting of 100 psychiatric patients—20 patients each of schizophrenia, depressive disorder, bipolar disorder, anxiety disorder and alcohol use disorder. & group 2 consisting of 100 medical patients.

Correspondence

Tanul Jain
PG Resident, Department of
Psychiatry, Index Medical
College, Indore,
Madhya Pradesh, India

Inclusion criteria

1. Male patients.
2. Ages from 18 to 65 years
3. Married and living with their wives
4. Written informed consent given

Exclusion criteria

1. Female patients.
2. Male patients who are single, widow, divorced, separated, or married but are not living with their wives.
3. Psychiatric patients with comorbid medical conditions.
4. Medical patients with comorbid psychiatric disorders.
5. Patients taking medications which are known to cause sexual dysfunction.
6. Male patients whose spouse is suffering from sexual dysfunction.
7. Not given written informed consent.

Diagnosis of psychiatric disorders was done according to the criteria of DSM-5. Diagnosis of other medical conditions was done by respective department. Data collection tools were demographic data questionnaire. All patients were subjected to IIEF (International Index of Erectile Function) Questionnaire. The statistical analysis of collected data was done by using the SPSS program v26 (Statistical Package of Social Science).

Results

- There was a positive relationship established between Psychiatric ill male patients and sexual dysfunction.
- Sexual dysfunction was significantly high in mentally ill patients i.e., 59% compared to our control group i.e.,

- other medical patients (23%).
- Among the Psychiatric disorders highest frequency was observed in Schizophrenia which was 73%. (Figure -1)
- Other prevalence was: bipolar disorders-49%, depressive disorders-51%, anxiety disorders – 36% and Alcohol Use Disorder-58%.
- Among our control group the maximum frequency was observed in patients with heart diseases which was 41%. Other prevalence were liver disorders-25%, Diabetic-20%, genitourinary tract disorders-15% and respiratory disorders -10%.
- Psychiatric patients had a statistically significant higher orgasmic dysfunction, sexual desire dysfunction, intercourse dissatisfaction, and overall sexual dissatisfaction than other medical patients (*p* value < 0.05), with no statistically significant difference (Table-3) between the patients of both groups as regards erectile dysfunction (*p* value > 0.05).
- Among the patients of group, I (psychiatric patients), patients with schizophrenia had a statistically significant higher orgasmic dysfunction, intercourse dissatisfaction, and overall sexual dissatisfaction than patients with alcohol use disorder, depressive disorders, bipolar disorders, and anxiety disorders respectively (*p* value < 0.05), with no statistically significant difference between them as regards erectile dysfunction and sexual desire dysfunction (*p* value > 0.05) (Table-1).
- Among the patients of group II (other medical patients), there was no statistically significant difference between them as regards erectile dysfunction, orgasmic dysfunction, sexual desire dysfunction, intercourse dissatisfaction, and overall sexual dissatisfaction (*p* value > 0.05) (Table-2).




Fig 1: Comparison of Prevalence of Sexual Dysfunction

Table 1: Patterns of sexual functions among patients of group I (psychiatric patients)

IEEF		Schizophrenia (n=20)	Bipolar Disorder (n=20)	Depression (n=20)	Anxiety (n=20)	Alcohol Use Disorder (n=20)
Erectile function	Normal	3	10	9	12	6
	Abnormal	17	10	11	8	14
Orgasmic function	Normal	5	11	11	11	7
	Abnormal	15	9	9	9	13
Sexual desire	Normal	7	10	7	13	11
	Abnormal	13	10	13	7	9
Intercourse satisfaction	Normal	7	13	10	12	10
	Abnormal	13	12	10	8	10
Overall satisfaction	Normal	4	12	12	16	8
	Abnormal	16	8	8	4	12

Table 2: Patterns of sexual functions among patients of group II (other medical patients)

IEEF		Cardiac Disease (n=20)	Liver disease (n=20)	Diabetes (n=20)	Genitourinary disease (n=20)	Respiratory Disorder (n=20)
Erectile function	Normal	10	13	14	16	17
	Abnormal	10	7	6	4	3
Orgasmic function	Normal	12	15	15	17	18
	Abnormal	8	5	5	3	2
Sexual desire	Normal	11	16	18	18	18
	Abnormal	9	4	2	2	2
Intercourse satisfaction	Normal	13	15	17	17	18
	Abnormal	7	5	3	3	2
Overall satisfaction	Normal	13	16	16	17	19
	Abnormal	7	4	4	3	1

Table 3: Patterns of sexual functions among patients of group I and group II

IEEF		Group I (n=100)	Group II (n=100)
Erectile function	Normal	40	70
	Abnormal	60	30
Orgasmic function	Normal	45	77
	Abnormal	55	23
Sexual desire	Normal	48	81
	Abnormal	52	19
Intercourse satisfaction	Normal	47	80
	Abnormal	53	20
Overall satisfaction	Normal	52	81
	Abnormal	48	19

Conclusion

Prevalence of sexual dysfunctions are higher among psychiatric ill male patients than medical patients therefore more attention should be given to the treatment of these disorders and counselling should also be done. The persistence of sexual problems has significant negative impact on patient’s satisfaction and adherence with the treatment, quality of life, and partnership. Routine assessment of sexual functioning needs to be integrated into ongoing care to identify and address problems early. If sexual dysfunction is ignored, it may maintain the psychiatric disorder, compromise treatment outcome, and lead to non-adherence.

Conflict of Interest

Not available

Financial Support

Not available

References

1. Steven A, Price J. Evolutionary psychiatry; A new beginning, 2nd edn. Routledge, London; c2000.
2. Rahmani A, Merghati E, AllahGholi L. Sexual satisfaction and its relation to marital happiness in Iranians. Iran J Publ Heal. 2012;38:77-82
3. Dunn KM, Croft PR, Halkett GI. Association of sexual problems with social, psychological and physical problems in men and women. Epidemiol Comm Heal. 1999;53:144-148
4. Bancroft JHJ. Human sexuality and its problems, 2nd edn. Churchill Livingstone, Edinburgh; c1989
5. Rowland DL, Incrocci L. Handbook of sexual and gender identity disorders. Hoboken, N.J.: John Wiley & Sons; c2008. p. 80.
6. Howes OD, Wheeler MJ, Pilowsky LS, Landau S, Murray RM, Smith S. Sexual function and gonadal hormones in patients taking antipsychotic treatment for schizophrenia or schizoaffective disorder. J Clin Psych. 2007;68:361-367
7. Montejo AL, Majadas S, Rico-Villademoros F, Lorca G, De La Gandara J, Franco M. Frequency of sexual dysfunction in patients with a psychotic disorder receiving antipsychotics. J Sex Med. 2010;7:3404-3413.
8. Westheide J, Cohen S, Bender S, Cooper-Mahkorn D, Erfurth A, Gastpar M. Sexual dysfunction in psychiatric inpatients the role of antipsychotic medication.

- Pharmacopsych. 2007;40:140-145.
9. Malik P, Kemmler G, Hummer M, Riecher-Rössler A, Kahn RS, Fleischhacker WW. Sexual dysfunction in first-episode schizophrenia patients: results from European first episode schizophrenia trial. *J Clin Psychopharmacol.* 2015;31:274-280.
 10. Rosen R, Riley A, Wagner G, Osterloh I, Kirkpatrick J, Mishra A. The international index of erectile function (IIEF): A multidimensional scale for assessment of erectile dysfunction. *Urol.* 1997;49:822-830.
 11. Shamloul R, Ghanem H, Abou-zeid A. Validity of the Arabic version of the sexual health inventory for men among Egyptians. *Int J Impot Res.* 2004;16(5):452-455.

How to Cite This Article

Jain T, Razdan RG. Study of prevalence of sexual dysfunction in males with psychiatric illness compared to other medical patients. *International Journal of Research in Psychiatry.* 2023;3(1):12-15.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.